Reach S-A80 (Temporary Access Road) Intermittent Spread D Webster County, West Virginia

| Data | Included |
|--|---------------|
| Photos | \checkmark |
| SWVM Form | \checkmark |
| FCI Calculator and HGM Form | \checkmark |
| RBP Physical Characteristics Form | \checkmark |
| Water Quality Data | N/A – No flow |
| RBP Habitat Form | \checkmark |
| RBP Benthic Form | \checkmark |
| Benthic Identification Sheet | N/A – No flow |
| Wolman Pebble Count | \checkmark |
| Reference Reach Software Pebble Count Data | \checkmark |
| Longitudinal Profile and Cross Sections | \checkmark |

Spread D Stream S-A80 (Temporary Access Road) Webster County



Photo Type: DS, US View Location, Orientation, Photographer Initials: Downstream at ROW looking E upstream, COC Lat: 38.480687 Long: -80.554061



Photo Type: DS, DS View Location, Orientation, Photographer Initials: Downstream at ROW looking SW downstream, COC Lat: 38.480687 Long: -80.554061

DEP Draft Permit #WQC-21-005

Spread D Stream S-A80 (Temporary Access Road) Webster County



Photo Type: CL US

Location, Orientation, Photographer Initials: On thalweg at ROW centerline looking SE Upstream, COC Lat: 38.480687 Long: -80.554061



Photo Type: CL DS Location, Orientation, Photographer Initials: On thalweg at ROW centerline looking W Downstream, COC Lat: 38.480687 Long: -80.554061

DEP Draft Permit #WQC-21-005

Spread D Stream S-A80 (Temporary Access Road) Webster County



Photo Type: US, US View Location, Orientation, Photographer Initials: Upstream at ROW looking SE upstream, COC Lat: 38.480687 Long: -80.554061



Photo Type: US, DS View Location, Orientation, Photographer Initials: Upstream at ROW looking SW downstream, COC Lat: 38.480687 Long: -80.554061

DEP Draft Permit #WQC-21-005

West Virginia Stream and Wetland Valuation Metric (SWVM) Version 2.1, September 2017

| USACE FILE NO./ Project Name: (v2.1, Sept 2015) | | Mountain | Valley Pipeline | | COORDINATES: cimal Degrees) | Lat. | 38.480687 | Lon. | -80.554061 | WEATHER: | Partly Cloudy | DATE: | 9/9/2021 |
|---|----------------------|------------------------|--|---------------------|--------------------------------|------|--|-----------------------|-------------|--|-------------------------------|--|------------------------------|
| IMPACT STREAM/SITE ID (watershed size (acreage). | | | Ş. | A80 | | | MITIGATION STREAM CLASS. (watershed size {acreag | | | | - | Comments: | |
| STREAM IMPACT LENGTH: | 104 | FORM OF MITIGATION: | RESTORATION (Levels I-III) | | OORDINATES: cimal Degrees) | Lat. | | Lon. | | PRECIPITATION PAST 48 HRS: | | Mitigation Length: | |
| Column No. 1- Impact Existing | g Condition (De | bit) | Column No. 2- Mitigation Existing C | ondition - Base | line (Credit) | | Column No. 3- Mitigation P Post Completio | | ears | Column No. 4- Mitigation Proje Post Completion (| | Column No. 5- Mitigation Projecte | d at Maturity (Credit) |
| Stream Classification: | Intern | nittent | Stream Classification: | | | | Stream Classification: | | 0 | Stream Classification: | 0 | Stream Classification: | 0 |
| Percent Stream Channel Sl | оре | 4.6 | Percent Stream Channel Sl | ope | | | Percent Stream Channel S | lope | 0 | Percent Stream Channel SI | ope 0 | Percent Stream Channel SI | ope 0 |
| HGM Score (attach d | ata forms): | | HGM Score (attach | data forms): | | | HGM Score (attach | data forms): | | HGM Score (attach da | ata forms): | HGM Score (attach da | ita forms): |
| | 0.51 | Average | | | Average | | | | Average | | Average | | Avera |
| łydrology Biogeochemical Cycling Habitat | 0.51 0.32 0.09 | 0.30666667 | Hydrology Biogeochemical Cycling Habitat | | 0 | | Hydrology Biogeochemical Cycling Habitat | | 0 | Hydrology Biogeochemical Cycling Habitat | 0 | Hydrology Biogeochemical Cycling Habitat | 0 |
| PART I - Physical, Chemical and | | cators | PART I - Physical, Chemical an | d Biological Ind | licators | | PART I - Physical, Chemical a | nd Biological Ind | icators | PART I - Physical, Chemical and | Biological Indicators | PART I - Physical, Chemical and | Biological Indicators |
| | Points Scale Range | Site Score | | Points Scale Range | Site Score | | | Points Scale Range | Site Score | | Points Scale Range Site Score | | Points Scale Range Site Scor |
| HYSICAL INDICATOR (Applies to all streams | s classifications) | | PHYSICAL INDICATOR (Applies to all streams | classifications) | | | PHYSICAL INDICATOR (Applies to all stream | s classifications) | | PHYSICAL INDICATOR (Applies to all streams | classifications) | PHYSICAL INDICATOR (Applies to all streams | classifications) |
| SEPA RBP (High Gradient Data Sheet) | | | USEPA RBP (Low Gradient Data Sheet) | | | | USEPA RBP (High Gradient Data Sheet) | | | USEPA RBP (High Gradient Data Sheet) | | USEPA RBP (High Gradient Data Sheet) | |
| Epifaunal Substrate/Available Cover Embeddedness | 0-20 | 0 | 1. Epifaunal Substrate/Available Cover 2. Pool Substrate Characterization | 0-20 | | | 1. Epifaunal Substrate/Available Cover 2. Embeddedness | 0-20 | | 1. Epifaunal Substrate/Available Cover 2. Embeddedness | 0-20 | 1. Epifaunal Substrate/Available Cover 2. Embeddedness | 0-20 |
| Velocity/ Depth Regime | 0-20 | 0 | 3. Pool Variability | 0-20 | | | 3. Velocity/ Depth Regime | 0-20 | | 3. Velocity/ Depth Regime | 0-20 | 3. Velocity/ Depth Regime | 0-20 |
| Sediment Deposition | 0-20 | 1 | 4. Sediment Deposition | 0-20 | | | 4. Sediment Deposition | 0-20 | | 4. Sediment Deposition | 0-20 | 4. Sediment Deposition | 0-20 |
| . Channel Flow Status | 0-20 0-1 | 0 | 5. Channel Flow Status | 0-20 0-1 | | | 5. Channel Flow Status | 0-20 0-1 | | 5. Channel Flow Status | 0-20 0-1 | 5. Channel Flow Status | 0-20 0-1 |
| . Channel Alteration | 0-20 | 2 | 6. Channel Alteration | 0-20 | | | 6. Channel Alteration | 0-20 | | 6. Channel Alteration | 0-20 | 6. Channel Alteration | 0-20 |
| Frequency of Riffles (or bends) | 0-20 | 0 | 7. Channel Sinuosity | 0-20 | | | 7. Frequency of Riffles (or bends) | 0-20 | | 7. Frequency of Riffles (or bends) | 0-20 | 7. Frequency of Riffles (or bends) | 0-20 |
| Bank Stability (LB & RB) | 0-20 | 6 | 8. Bank Stability (LB & RB) | 0-20 | | | 8. Bank Stability (LB & RB) | 0-20 | | 8. Bank Stability (LB & RB) | 0-20 | 8. Bank Stability (LB & RB) | 0-20 |
| Vegetative Protection (LB & RB) | 0-20 | 6 | 9. Vegetative Protection (LB & RB) 10. Riparian Vegetative Zone Width (LB & RB) | 0-20 | | | 9. Vegetative Protection (LB & RB) 10. Riparian Vegetative Zone Width (LB & RB) | 0-20 | | 9. Vegetative Protection (LB & RB) 10. Riparian Vegetative Zone Width (LB & RB) | 0-20 | 9. Vegetative Protection (LB & RB) 10. Riparian Vegetative Zone Width (LB & RB) | 0-20 |
| 0. Riparian Vegetative Zone Width (LB & RB) otal RBP Score | 0-20 Poor | 29 | Total RBP Score | 0-20 Poor | 0 | | Total RBP Score | 0-20 Poor | 0 | Total RBP Score | 0-20 Poor 0 | Total RBP Score | Poor C |
| ub-Total | POOI | 0.145 | Sub-Total | POOL | 0 | | Sub-Total | POOL | 0 | Sub-Total | 0 | Sub-Total | 9001 0 |
| HEMICAL INDICATOR (Applies to Intermitten | nt and Perennial Str | | CHEMICAL INDICATOR (Applies to Intermittent | and Perennial Stre | eams) | | CHEMICAL INDICATOR (Applies to Intermitte | nt and Perennial Stre | ams) | CHEMICAL INDICATOR (Applies to Intermitten | | CHEMICAL INDICATOR (Applies to Intermitten | |
| VDEP Water Quality Indicators (General |) | | WVDEP Water Quality Indicators (General) | | | | WVDEP Water Quality Indicators (General |) | | WVDEP Water Quality Indicators (General |) | WVDEP Water Quality Indicators (General) | |
| pecific Conductivity | | | Specific Conductivity | | | | Specific Conductivity | | | Specific Conductivity | | Specific Conductivity | |
| 100-199 - 85 points | 0-90 | | | 0-90 | | | | 0-90 | | | 0-90 | | 0-90 |
| Ħ | 0.1 | | рн | 0.1 | | | рн | 0.1 | | рн | 0.1 | рн | 0,1 |
| 5.6-5.9 = 45 points | 0-80 | | | 5-90 | | | | 5-90 | | | 5-90 | | 5-90 |
| 0 | | | 00 | | | | DO | | | DO | | DO | |
| | 10-30 | | | 10-30 | | | | 10-30 | | | 10-30 | | 10-30 |
| | 10-00 | | l | 10-00 | | | | 10-30 | | | | | |
| ub-Total | | | Sub-Total | | 0 | | Sub-Total | | 0 | Sub-Total | 0 | Sub-Total | 0 |
| IOLOGICAL INDICATOR (Applies to Intermitt | tent and Perennial | Streams) | BIOLOGICAL INDICATOR (Applies to Intermitte | ent and Perennial S | Streams) | | BIOLOGICAL INDICATOR (Applies to Interr | nittent and Perenni | al Streams) | BIOLOGICAL INDICATOR (Applies to Interm | ittent and Perennial Streams) | BIOLOGICAL INDICATOR (Applies to Intermi | ttent and Perennial Stream |
| V Stream Condition Index (WVSCI) | 0.100 0.1 | | WV Stream Condition Index (WVSCI) | | | | WV Stream Condition Index (WVSCI) | | | WV Stream Condition Index (WVSCI) | | WV Stream Condition Index (WVSCI) | |
| 0 | 0-100 0-1 | | | 0-100 0-1 | | | | 0-100 0-1 | | | 0-100 0-1 | | 0-100 0-1 |
| Sub-Total | | 0 | Sub-Total | | 0 | | Sub-Total | | 0 | Sub-Total | 0 | Sub-Total | 0 |
| PART II - Index and U | Jnit Score | | PART II - Index and | Unit Score | | | PART II - Index an | d Unit Score | | PART II - Index and U | nit Score | PART II - Index and U | nit Score |
| | | | | | | | | | | | | | |
| Index | Linear Feet | Unit Score | Index | Linear Feet | Unit Score | | Index | Linear Feet | Unit Score | Index | Linear Feet Unit Score | Index | Linear Feet Unit Sc |

104 40.5166667

0.390

FCI Calculator for the High-Gradient Headwater Streams in Appalachia

To ensure accurate calculations, the UPPERMOST STRATUM of the plant community is determined based on the calculated value for V_{CCANOPY} (≥20% cover is required for tree/sapling strata). Go to the SAR Data Entry tab and enter site characteristics and data in the yellow cells. For information on determining how to split a project into SARs, see Chapter 5 of the Operational Draft Regional Guidebook for the Functional Assessment of High-Gradient Headwater Streams and Low-Gradient Perennial Streams in Appalachia (Environmental Laboratory U.S. Army Corps of Engineers 2017).

Project Name: MVP Location: Webster, Spread D Sampling Date: 9/9/21

Subclass for this SAR:

Intermittent Stream

Uppermost stratum present at this SAR: Shrub/Herb Strata

SAR number:

S-A80

Before Project

Functional Results Summary:

Enter Results in Section A of the Mitigation Sufficiency Calculator

Project Site

| Function | Functional Capacity Index |
|------------------------|------------------------------|
| Hydrology | 0.51 |
| Biogeochemical Cycling | 0.32 |
| Habitat | 0.09 |

Variable Measure and Subindex Summary:

| Variable | Name | Average Measure | Subindex |
|------------------------|---|--------------------|----------|
| VCCANOPY | Percent canpoy over channel. | Not Used, <20% | Not Used |
| VEMBED | Average embeddedness of channel. | 1.70 | 0.35 |
| V _{SUBSTRATE} | Median stream channel substrate particle size. | 0.08 | 0.04 |
| V _{BERO} | Total percent of eroded stream channel bank. | 4.08 | 1.00 |
| V _{LWD} | Number of down woody stems per 100 feet of stream. | 1.02 | 0.13 |
| V _{TDBH} | Average dbh of trees. | Not Used | Not Used |
| V _{SNAG} | Number of snags per 100 feet of stream. | 0.00 | 0.10 |
| V _{SSD} | Number of saplings and shrubs per 100 feet of stream. | 25.51 | 0.39 |
| V _{SRICH} | Riparian vegetation species richness. | 0.00 | 0.00 |
| V _{DETRITUS} | Average percent cover of leaves, sticks, etc. | 6.88 | 0.08 |
| V _{HERB} | Average percent cover of herbaceous vegetation. | 20.00 | 0.27 |
| V _{WLUSE} | Weighted Average of Runoff Score for Catchment. | 0.89 | 0.94 |

| | | | Field D | Data She | et and C | alculato | r | | | |
|------------------------|---|------------------------------|--------------------------|--------------------------|------------------------------|--------------------------|--------------------------|---------------------------------|-------------|-----------------|
| Team | RFC, COC | | | | | | Latitude/UT | M Northing: | 38.480687 | |
| Project Name | MVP | | | | | L | .ongitude/U | TM Easting: | -89.554061 | |
| Location | Webster, S | pread D | | | | | San | npling Date: | 9/9/21 | |
| SAR Number | S-A80 | Reach | Length (ft): | 98 | Stream Ty | ype: Inter | mittent Strea | im | | |
| Top Strata | Sh | rub/Herb Sti | ata | (determine | d from perce | ent calculate | ed in V _{CCANO} | _{PY}) | | |
| Site and Timing | 10.04 | 8 | | | • | Before Proje | ct | | | • |
| nple Variables | 1-4 in strea Average pe | | | al hu traa au | d conling c | anany Mar | aura at na f | ower then 1 | 0 roughly | |
| 1 V _{CCANOPY} | equidistant 20%, enter | points alonç at least one | the stream value betw | . Measure een 0 and 1 | | apling cove | r is at least | 20%. (If les | | Not Use <20% |
| List the pe | rcent cover r | lleasureillei | its at each | Joint Delow. | | | | | | |
| | | | | | | | | | | |
| 2 V _{EMBED} | | | | | | | | ghly equidis | | 1.7 |
| | | | | | | | | percentage | | 1.7 |
| | to the follow | ving table. I | f the bed is | an artificial s | | composed o | | ter the rating ents, use a r | | |
| | - | ness rating | | | • | | d from Plat | s, Megahan | , and | |
| | Rating | Rating Des | cription | | | | | | | |
| | 5 | <5 percent | of surface c | | | | | (or bedrock | () | |
| | 4 | | | | | | | | | |
| | 3 | | | | d, surrounde d, surrounde | | | | | |
| | 1 | | | | | | | nt (or artificia | al surface) | |
| | ings at each | | | | | | | | | |
| 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 5 | 1 | |
| 1 | 1 | 3 | 5 | 4 | 1 5 | 1 | 1 | 2 | 1 | |
| 1 | 1 | | | 4 | 5 | 1 | | 1 | | |
| | | | | | | | | | | |
| | Median stre along the s cle size in ine | tream; use t | he same po | ints and par | ticles as use | ed in V _{EMBED} |). | | | 0.08 i |
| | e as 0.0 in, s | | | | | `` | | | | |
| 0.08 | 0.08 | 0.08 | 0.08 | 0.40 | 0.08 | 0.08 | 0.08 | 1.50 | 0.08 | |
| 0.08 | 0.08 | 0.60 0.08 | 1.30 0.08 | 0.70 | 0.08 | 0.08 | 0.80 | 0.70 | 0.08 | |
| 0.08 | 0.08 | 0.08 | 0.08 | 1.20 | 0.00 | 0.06 | 0.08 | 0.06 | 0.08 | |
| 4 V _{RERO} | Total perce | nt of eroder | stream cha | nnel bank | Enter the to | tal number | of feet of er | oded bank o | n each side | |
| 4 V _{BERO} | and the tota | al percentag | | | | | | for the stream | | 4 % |
| | up to 200% | Left Bank: | 2 | ft | | Right Bank: | 2 | : ft | | |
| nple Variables | 5-9 within t | he entire rij | oarian/buff | er zone adia | acent to the | stream ch | annel (25 fe | et from ea | ch bank). | |
| 5 V _{LWD} | | | | - | | | • | th) per 100 | - | |
| | | ch. Enter th t of stream | | | e 50'-wide b | ouffer and w | ithin the cha | innel, and th | e amount | 1.0 |
| | | | | | downed wo | ody stems: | | 1 | | |
| 6 V _{tdbh} | | | | | | ig cover is a | t least 20%) | . Trees are | at least 4 | Not Us |
| | | , | | tree DBHs i | | | | | | |
| | List the dbh the stream | | ents of indiv | idual trees | at least 4 in | i) within the | buffer on ea | ich side of | | |
| | Sie er ourit | Left Side | | | | | Right Side | | | |
| 16 | | | | | 0 | | _ | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |
| 7 V _{SNAG} | | | | | | | Enter numb | er of snags | on each | |
| | side of the | stream, and | the amount | per 100 fee | et will be cal | culated. | | | | 0.0 |
| | | Left Side: | | 0 | | Right Side: | | 0 | | |
| B V _{SSD} | Number of | | | - | up to 4 inch | | | stream (me | asure only | |
| | if tree cove | r is <20%). | Enter numb | er of sapling | gs and shrul | bs on each : | side of the s | tream, and t | the amount | 25.5 |
| | | f stream wil | ho ocloui- | od | | | | | | |

| 9 | V _{SRICH} | Group 1 in | the tallest st | tratum. Ch | eck all exotic | and invas | am reach. Ch ive species p from these d | resent in all | | | 0.00 |
|------|---|-----------------------------------|-----------------------------------|-------------------------------|------------------------------|------------------------------------|--|------------------------------|--------------|----------------|-----------------------|
| | | | 0.1 = 1.0 | | | | | | 2 (-1.0) | | |
| | Acer rubru | | | Magnolia t | ripetala | | Ailanthus a | <u> </u> | | Lonicera ja | oonica |
| | Acer sacch | | | Nyssa sylv | - | | Albizia julib | | | Lonicera ta | |
| | Aesculus fi | | | | n arboreum | | Alliaria peti | | | Lotus corni | |
| | Asimina tril | | | Prunus sei | | | | | | Lythrum sa | |
| | Betula alleg | | | Quercus a | | | Alternanthe philoxeroide | | | Microstegium | |
| | - | | | | | | | | | Paulownia | |
| | Betula lent | | | Quercus c | | | Aster tatari | | | | |
| | Carya alba | | | Quercus in | | | Cerastium | | | Polygonum o | |
| | Carya glab | | | Quercus p | | | Coronilla va | | | Pueraria m | |
| | Carya oval | | | Quercus ru | | | Elaeagnus u | | | Rosa multit | |
| | Carya ovat | a | | Quercus v | elutina | | Lespedeza | bicolor | | Sorghum h | |
| | Cornus flor | rida | | Sassafras | albidum | | Lespedeza | cuneata | | Verbena br | asiliensis |
| | Fagus grar | ndifolia | | Tilia ameri | cana | | Ligustrum ot | otusifolium | | | |
| | Fraxinus a | mericana | | Tsuga can | adensis | | Ligustrum s | sinense | | | |
| | Liriodendron | tulipifera | | Ulmus am | ericana | | | | | | |
| | Magnolia a | cuminata | | | | | | | | | |
| _ | | | | <u> </u> | | | | | | | |
| Comm | la Variablea | 0 | Species in | | 40" × 40" o | r 4m v 4m |) in the rineri | 0 | Species in | • | aaah |
| | | b plots shou Average pe | IId be place | d roughly of leaves, | equidistant | ly along e a ier organic |) in the ripari ach side of the material. Wo ayer at each s | ne stream. body debris | | | 6.88 % |
| | | | Left | Side | | | Right | Side | |] | |
| | | 0 | 0 | 0 | 0 | 30 | 20 | 0 | 5 | | |
| | | | | | | | | | | | |
| 11 | V _{HERB} | include woo | ody stems a percentages ot. | t least 4" dt s up througl | oh and 36" ta | all. Becaus | e there may b Enter the per | e several la cent cover c | yers of grou | and cover | 20 % |
| | | | | Side | - | 70 | | Side | 0 | | |
| | | 0 | 0 | 0 | 5 | 70 | 80 | 5 | 0 | | |
| 12 | V _{wluse} | Weighted A | | | e for watersh se From Dro | | | | Runoff | % in Catch- | 0.89 Running |
| | French and a | atter concerts | | | | p List) | | _ | Score | ment | Percent (not >100) |
| Î | | ative range (: | | ASSA KUR | | | | • | 1 | 87.3 | 87.3 |
| | Open space | (pasture, law | ns, parks, etc.) |), grass cover | <50% | | | • | 0.1 | 4.3 | 91.6 |
| | Impervious | areas (parking | lots, roofs, d | riveways, etc | :) | | | - | 0 | 3.9 | 95.5 |
| | Open space | (pasture, law | ns, parks, etc. |), grass cover | 50% - 75% | | | - | 0.2 | 1.61 | 97.11 |
| Î | | | | | | | | | | | |
| | - | (pasture, law | · | | -1076 | | | | 0.3 | 1.38 | 98.49 |
| | Residential of | districts, 2 acro | es (12% cover |) | | | | • | 0.3 | 1.51 | 100 |
| Î | | | | | | | | - | | | |
| Î | | | | | | | | - | | | |
| | - | | | | | | •• | | | | |
| | | S-A80 | | | | | No | tes: | | | |
| ١ | /ariable | Value | VSI | | | | | | | | |
| | VCCANOPY | Not Used, <20% | Not Used | | | | | | | | |
| | VEMBED | 1.7 | 0.35 | | | | | | | | |
| | V _{SUBSTRATE} | 0.08 in | 0.04 | | | | | | | | |
| , | V _{BERO} | 4 % | 1.00 | | | | | | | | |
| , | | 1.0 | 0.13 | | | | | | | | |
| , | V _{TDBH} | Not Used | Not Used | | | | | | | | |
| | V _{SNAG} | 0.0 | 0.10 | | | | | | | | |
| ļ , | | 25.5 | | | | | | | | | |
| | V _{SSD} | 25.5 | 0.39 | | | | | | | | |
| | | | 0.39 | | | | | | | | |
| | V _{SRICH} | 0.00 | 0.00 | | | | | | | | |
| , | V _{SRICH} V _{DETRITUS} | 0.00 6.9 % | 0.00 0.08 | | | | | | | | |
| , | V _{SRICH} | 0.00 | 0.00 | | | | | | | | |

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

| STREAM NAME | LOCATION | | | | |
|---------------------|---------------|-------------------|--|--|--|
| STATION # RIVERMILE | STREAM CLASS | | | | |
| LAT LONG | RIVER BASIN | | | | |
| STORET # | AGENCY | | | | |
| INVESTIGATORS | INVESTIGATORS | | | | |
| FORM COMPLETED BY | DATE TIME | REASON FOR SURVEY | | | |

| | Has there been a beauty rain in the last 7 days? |
|----------------------------|---|
| WEATHER CONDITIONS | Now Past 24 hours Has there been a heavy rain in the last 7 days? Storm (heavy rain) rain (steady rain) showers (intermittent) Yes No % %cloud cover clear/sunny Mir Temperature0 C |
| SITE LOCATION/MAP | Draw a map of the site and indicate the areas sampled (or attach a photograph) |
| | Access Road |
| STREAM CHARACTERIZATION | Stream Subsystem Perennial Tidal Stream Type Coldwater Warnwater Stream Origin Glacial Spring-fed Mixture of origins Catchment Area km² Swamp and bog Other Other Km² |

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

| WATERSHED FEATURES RIPARIAN VEGETATION (18 meter buffer) | Predominant Surrounding Landuse Local Watershed NPS Pollution Forest Commercial Field/Pasture Industrial Agricultural Other Residential Other Indicate the dominant type and record the dominant species present Herbaceous Trees Shrubs Grasses Dominant species present Herbaceous |
|--|--|
| INSTREAM FEATURES | Dominant species present |
| LARGE WOODY | LWDm ² |
| DEBRIS | Density of LWDm ² /km ² (LWD/ reach area) |
| AQUATIC | Indicate the dominant type and record the dominant species present |
| VEGETATION | Rooted emergent Rooted submergent Rooted floating Free floating Floating Algae Attached Algae Booted floating Free floating Free floating Dominant species present |
| WATER QUALITY (DS, US) | Temperature0 C Water Odors Normal/None Sewage Specific Conductance Petroleum Fishy Chemical Other Dissolved Oxygen Water Surface Oils Slick Sheen None Globs Flecks pH Turbidity (if not measured) Clear Slightly turbid Turbid Turbid Turbid Opaque Turbid |
| SEDIMENT/ | Odors |
| SUBSTRATE | Normal Sewage Petroleum Deposits Chemical Anaerobic None Sludge Sawdust Paper fiber Sand Other Other Epoking at stones which are not deeply embedded are the undersides black in color? How are the undersides black in color? |

| INC | ORGANIC SUBSTRATE (should add up to | | ORGANIC SUBSTRATE COMPONENTS (does not necessarily add up to 100%) | | | | |
|-------------------|--|------------------------------------|---|----------------------------|-----------------------------------|--|--|
| Substrate Type | Diameter | % Composition in Sampling Reach | Substrate Type | Characteristic | % Composition in Sampling Area | | |
| Bedrock | | | Detritus | sticks, wood, coarse plant | | | |
| Boulder | > 256 mm (10") | | | materials (CPOM) | | | |
| Cobble | 64-256 mm (2.5"-10") | | Muck-Mud | black, very fine organic | | | |
| Gravel | 2-64 mm (0.1"-2.5") | | | (FPOM) | | | |
| Sand | 0.06-2mm (gritty) | | Marl | grey, shell fragments | | | |
| Silt | 0.004-0.06 mm | | | | | | |
| Clay | < 0.004 mm (slick) | | | | | | |

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

| STREAM NAME | LOCATION | | | | |
|---------------------|--------------------|-------------------|--|--|--|
| STATION # RIVERMILE | STREAM CLASS | | | | |
| LAT LONG | RIVER BASIN | | | | |
| STORET # | AGENCY | | | | |
| INVESTIGATORS | INVESTIGATORS | | | | |
| FORM COMPLETED BY | DATE TIME AM PM | REASON FOR SURVEY | | | |

| | Habitat | | Condition | ı Category | |
|--|---|---|---|---|---|
| | Parameter | Optimal | Suboptimal | Marginal | Poor |
| | 1. Epifaunal Substrate/ Available Cover | Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are <u>not</u> new fall and <u>not</u> transient). | 40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale). | 20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed. | Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking. |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |
| n sampling reach | 2. Embeddedness | Gravel, cobble, and boulder particles are 0- 25% surrounded by fine sediment. Layering of cobble provides diversity of niche space. | Gravel, cobble, and boulder particles are 25- 50% surrounded by fine sediment. | Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment. | Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment. |
| ted i | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |
| Parameters to be evaluated in sampling reach | 3. Velocity/Depth Regime | All four velocity/depth regimes present (slow- deep, slow-shallow, fast- deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.) | Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes). | Only 2 of the 4 habitat regimes present (if fast- shallow or slow-shallow are missing, score low). | Dominated by 1 velocity/ depth regime (usually slow-deep). |
| uram | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |
| Pa | 4. Sediment Deposition | Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition. | Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools. | Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent. | Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition. |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |
| | 5. Channel Flow Status | Water reaches base of both lower banks, and minimal amount of channel substrate is exposed. | Water fills >75% of the available channel; or <25% of channel substrate is exposed. | Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed. | Very little water in channel and mostly present as standing pools. |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |

Rapid Bioassessment Protocols For Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates, and Fish, Second Edition - Form 2

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

| Habitat | | Condition | 1 Category | | | | | |
|--|--|--|--|---|--|--|--|--|
| Parameter | Optimal | Suboptimal | Marginal | Poor | | | | |
| 6. Channel Alteration | Channelization or dredging absent or minimal; stream with normal pattern. | Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present. | Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted. | Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely. | | | | |
| SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | | |
| 7. Frequency of Riffles (or bends) | Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important. | Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15. | Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25. | Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25. | | | | |
| SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | | |
| SCORE 8. Bank Stability (score each bank) Note: determine left or right side by facing downstream. SCORE (LB) SCORE (RB) 9. Vegetative Protection (score each bank) | Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected. | Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion. | Moderately unstable; 30- 60% of bank in reach has areas of erosion; high erosion potential during floods. | Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars. | | | | |
| SCORE (LB) | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | |
| SCORE (RB) | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | |
| 9. Vegetative Protection (score each bank) | More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally. | 70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well- represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining. | 50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining. | Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height. | | | | |
| SCORE(LB) | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | |
| SCORE(RB) | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | |
| 10. Riparian Vegetative Zone Width (score each bank riparian zone) | Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone. | Width of riparian zone 12-18 meters; human activities have impacted zone only minimally. | Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal. | Width of riparian zone <6 meters: little or no riparian vegetation due to human activities. | | | | |
| SCORE (LB) | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | |
| SCORE (RB) | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | |

Total Score _____

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

| STREAM NAME | | LOCATION | | | | | | |
|----------------------|--|--|-------------------|--|--|--|--|--|
| STATION # | _ RIVERMILE | STREAM CLASS | | | | | | |
| LAT | LONG | RIVER BASIN | | | | | | |
| STORET # | | AGENCY | AGENCY | | | | | |
| INVESTIGATORS | | | LOT NUMBER | | | | | |
| FORM COMPLETED | BY | DATE TIME | REASON FOR SURVEY | | | | | |
| HABITAT TYPES | Indicate the percentage of each habitat type present Cobble% Snags% Vegetated Banks% Sand% Submerged Macrophytes% Other ()% | | | | | | | |
| SAMPLE COLLECTION | Indicate the number of jab | lected? wading fi ps/kicks taken in each habitat ty lags Vegetated B | anks Sand | | | | | |
| GENERAL COMMENTS | | | | | | | | |

QUALITATIVE LISTING OF AQUATIC BIOTA

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare, 2 = Common, 3= Abundant, 4 = Dominant

| Periphyton | 0 | 1 | 2 | 3 | 4 | Slimes | 0 | 1 | 2 | 3 | 4 |
|-------------------|---|---|---|---|---|--------------------|---|---|---|---|---|
| Filamentous Algae | 0 | 1 | 2 | 3 | 4 | Macroinvertebrates | 0 | 1 | 2 | 3 | 4 |
| Macrophytes | 0 | 1 | 2 | 3 | 4 | Fish | 0 | 1 | 2 | 3 | 4 |

FIELD OBSERVATIONS OF MACROBENTHOS

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare (1-3 organisms), 2 = Common (3-9 organisms), 3= Abundant (>10 organisms), 4 = Dominant (>50 organisms)

| Porifera | 0 | 1 | 2 | 3 | 4 | Anisoptera | 0 | 1 | 2 | 3 | 4 | Chironomidae | 0 | 1 | 2 | 3 | 4 |
|-----------------|---|---|---|---|---|-------------|---|---|---|---|---|---------------|---|---|---|---|---|
| Hydrozoa | 0 | 1 | 2 | 3 | 4 | Zygoptera | 0 | 1 | 2 | 3 | 4 | Ephemeroptera | 0 | 1 | 2 | 3 | 4 |
| Platyhelminthes | 0 | 1 | 2 | 3 | 4 | Hemiptera | 0 | 1 | 2 | 3 | 4 | Trichoptera | 0 | 1 | 2 | 3 | 4 |
| Turbellaria | 0 | 1 | 2 | 3 | 4 | Coleoptera | 0 | 1 | 2 | 3 | 4 | Other | 0 | 1 | 2 | 3 | 4 |
| Hirudinea | 0 | 1 | 2 | 3 | 4 | Lepidoptera | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Oligochaeta | 0 | 1 | 2 | 3 | 4 | Sialidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Isopoda | 0 | 1 | 2 | 3 | 4 | Corydalidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Amphipoda | 0 | 1 | 2 | 3 | 4 | Tipulidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Decapoda | 0 | 1 | 2 | 3 | 4 | Empididae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Gastropoda | 0 | 1 | 2 | 3 | 4 | Simuliidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Bivalvia | 0 | 1 | 2 | 3 | 4 | Tabinidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| | | | | | | Culcidae | 0 | 1 | 2 | 3 | 4 | | | | | | |

WOLMAN PEBBLE COUNT FORM

| County: | Webster | Stream ID: | S-A80 |
|--------------|-----------------------|---------------|---------|
| Stream Name: | UNT to Laurel Creek | | |
| HUC Code: | | Basin: | |
| Survey Date: | 9/9/2021 | | |
| Surveyors: | RFC COC | Impact Reach: | 29.87 m |
| Type: | Representative/Riffle | | |

| | | | LE COUNT | | | | |
|-------------|--------------|-------------|---------------|-------------------|---------|--------|--------|
| Inches | PARTICLE | Millimeters | | Particle Count | Total # | Item % | % Cum |
| | Silt/Clay | < .062 | S/C | • | 62 | 62.00 | 62.00 |
| | Very Fine | .062125 | | • | 0 | 0.00 | 62.00 |
| | Fine | .12525 | | ▲ ▼ | 0 | 0.00 | 62.00 |
| | Medium | .255 | S A N D | ▲ ▼ | 0 | 0.00 | 62.00 |
| | Coarse | .50-1.0 | | ▲ ▼ | 0 | 0.00 | 62.00 |
| .0408 | Very Coarse | 1.0-2 | | ▲ ▼ | 0 | 0.00 | 62.00 |
| .0816 | Very Fine | 2 -4 | | • | 0 | 0.00 | 62.00 |
| .1622 | Fine | 4 -5.7 |] | ▲ ▼ | 0 | 0.00 | 62.00 |
| .2231 | Fine | 5.7 - 8 | | ▲ ▼ | 0 | 0.00 | 62.00 |
| .3144 | Medium | 8 -11.3 | | ▲ ▼ | 2 | 2.00 | 64.00 |
| .4463 | Medium | 11.3 - 16 | G R A V E L | * * | 2 | 2.00 | 66.00 |
| .6389 | Coarse | 16 -22.6 | | ▲ ▼ | 6 | 6.00 | 72.00 |
| .89 - 1.26 | Coarse | 22.6 - 32 | | ▲ ▼ | 22 | 22.00 | 94.00 |
| 1.26 - 1.77 | Vry Coarse | 32 - 45 | | ▲ ▼ | 4 | 4.00 | 98.00 |
| 1.77 -2.5 | Vry Coarse | 45 - 64 | | ▲ ▼ | 0 | 0.00 | 98.00 |
| 2.5 - 3.5 | Small | 64 - 90 | | ▲ ▼ | 2 | 2.00 | 100.00 |
| 3.5 - 5.0 | Small | 90 - 128 | COBBLE | ▲ ▼ | 0 | 0.00 | 100.00 |
| 5.0 - 7.1 | Large | 128 - 180 | COBBEE | ▲ ▼ | 0 | 0.00 | 100.00 |
| 7.1 - 10.1 | Large | 180 - 256 | | ▲ ▼ | 0 | 0.00 | 100.00 |
| 10.1 - 14.3 | Small | 256 - 362 | | ▲ ▼ | 0 | 0.00 | 100.00 |
| 14.3 - 20 | Small | 362 - 512 | | ▲ ▼ | 0 | 0.00 | 100.00 |
| 20 - 40 | Medium | 512 - 1024 | B O U L D E R | ▲ ▼ | 0 | 0.00 | 100.00 |
| 40 - 80 | Large | 1024 -2048 | | ▲ ▼ | 0 | 0.00 | 100.00 |
| 80 - 160 | Vry Large | 2048 -4096 | | ▲ ▼ | 0 | 0.00 | 100.00 |
| | Bedrock | | BDRK | ▲ ▼ | 0 | 0.00 | 100.00 |
| | | | | Totals: | 100 | | |
| | Total Tally: | | | | | | |



boulder

0%

Bankfull Channel Pebble Count, S-A80

D84

D95

27

35





- 1720

1714

- 1712

- 1710

SURVEY NOTES:

- 1. THIS MAP HAS BEEN ORIENTED TO NAD 1983 UTM ZONE 17N, AND VERTICALLY TO THE NORTH AMERICAN VERTICAL DATUM OF 1988 (NAVD 88), USING REAL TIME DGPS. FIELD LOCATIONS WERE COMPLETED ON SEPTEMBER 9, 2021.
- 2. EASEMENT LINES SHOWN ON PLAN VIEW WERE PROVIDED BY MOUNTAIN VALLEY PIPELINE.
- 3. SURVEY POINTS FOR CROSS SECTIONS AND THALWEG PROFILES COLLECTED IN 2021 HAVE BEEN USED IN COMBINATION WITH SURVEY POINTS COLLECTED PREVIOUSLY IN 2020 IN ORDER TO GENERATE THE PRE-CROSSING SURFACE SHOWN IN PLAN. DUE TO NATURAL EROSIONAL STREAM PROCESSES THAT CAN OCCUR OVER TIME, MINOR ADJUSTMENTS TO THE PROFILE ALIGNMENTS MAY HAVE BEEN REQUIRED IN ORDER TO GENERATE A CLEAN PRE-CROSSING SURFACE.
- 4. ALL SECTION VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.
- 5. POST-CROSSING SURVEY INFORMATION SHOWN IN RED. DATA PENDING.
- 6. POST-CROSSING SURVEY POINTS FOR CROSS SECTIONS AND THALWEG ARE PROJECTED ONTO PRE-CROSSING SECTION AND PROFILE VIEWS FOR COMPARISON.



